

Behavioral Genetics: Concepts and Challenges

[See online here](#)

The notion that human behavior could be affected and predetermined by genetic basis started to emerge in the 20th century. Unfortunately, during that time, the eugenics movement was associated with the emergence of this theory, which played a part in discrediting it. Recently, several twin studies showed similar behavioral tendencies in monozygotic twins, even if separated at a very early age, again signifying the role of behavioral genetics. Our current understanding of genome-wide associations with different behavioral, psychiatric and medical problems allowed us to confirm that behavioral genetics is a true science and behavior can be affected by genetic, biologic and molecular determinants.



Twin and Adoption Studies and Behavioral Genetics

The behavior of children who are adopted since birth is expected to resemble their adoptive parents if the environment plays the most important role. Surprisingly, early studies in behavioral genetics found that different behavioral traits in adopted children resemble their biologic and not their adoptive parents.

Intellectual ability, alcoholism and criminal behavior of the offspring clearly resembled their biologic parents in these studies. Additionally, the risk of **mental disorders**, such as mood disorders and **schizophrenia** was **similar to the risk in the biologic parents' families** and not the adoptive family.

Even more surprisingly, adopted children who were raised by schizophrenic parents or by parents with mental disorders were not at an increased risk of developing a mental disease, further emphasizing the importance of genetic predisposition to these disorders.

The next step in confirming the role of genetics in determining behavioral traits was to study twins. **Monozygotic twins** are thought to be genetically identical, while dizygotic twins share only 50 % of their genetic makeup.

Twin studies focusing on identifying a **positive association between certain traits** hypothesize that the similarities in behavioral traits should be higher in monozygotic twins than dizygotic twins.

Interestingly, personality traits, social attitudes, and even divorce rates were more similar in monozygotic twins than dizygotic ones, confirming the importance of genetics in affecting behavior. **Monozygotic concordance studies** never reach a perfect score of 1, but **usually, yield a ratio that is above 0.5** when there is a genetic association. This finding shows that the environment still plays an important role in determining the strength of expression of the predetermined behavioral traits.


Another approach to twin studies was to investigate the emergence of certain behavioral traits of twins who are raised apart. Again, the findings of these studies showed similarities between monozygotic twins who are raised apart, but the resemblance was not perfect, emphasizing the importance of the environment as well.

Behavioral Genetics and Development

The next logical step in twin and adoption studies was to study the **role of age and development** on genetic predisposing to certain behavioral traits.

These studies showed clearly that children's behavior is strongly determined by their environment rather than genetics, while adults' mental health, antisocial behavior, and social attitudes are largely based on genetic predisposition rather than the environment. For instance, schizophrenia, a thought disorder, was found to be highly associated with certain genes and to be little related to the environment.

Genetic-Environment Interplay



“Interaction between Heredity and Environmental Influences”
Image created by Lecturio

Because the risk of developing certain behavioral traits in adopted children and monozygotic twins was never solely explained by genetic predisposition, psychologists reached a hypothesis that **genetic predisposition** puts the subject at an increased risk of developing certain traits, and the **environment then can either ameliorate or increase** the strength of these traits.

This genetic–environment interplay has become of tremendous interest and importance to behavioral genetic scientists. In simple words, they hypothesize that genetics contribute towards the similarity between related individuals, while the environment is responsible for the differences.

The effect of genetics on overall IQ is challenging to understand for several reasons. Firstly, children’s IQ is related to their environment, their upbringing and is little affected by genetic predisposition unless a known neurodevelopmental disease is evident.

Secondly, the ultimate IQ of adults seems to be genetically determined rather than affected by the environment, emphasizing the role of development and age on behavioral genetics.

Finally, monozygotic twins and hereditary studies of IQ failed to reach a concordance level above 0.5, emphasizing that genetics alone cannot explain IQ, and **genetic–environment interplay is a real thing and an important determinant of IQ.**

Genome-Wide Association Studies and Behavioral Genetics

Neuroscientists and geneticists recently started to focus on **genetic polymorphisms** and their association with different biologic, physiologic, mental and behavioral traits. Genetic polymorphism is defined as

1. The existence together of **many forms of DNA sequences at a locus within the population.**
2. A discontinuous **genetic** variation that results in **different forms or types of individuals** among the members of a single species.

This recent work allowed the identification of several genetic loci where several

polymorphisms can occur and each of them is associated with an increased risk of certain traits.

It is important to note that a single polymorphism rarely has a causative relationship with a certain trait. However, the **interplay between several polymorphisms in different loci** eventually contributes to a phenotype that either has a positive or a negative trait. This trait can be behavioral, mental or physical.

One of the earliest polymorphisms to be studied in behavioral genetics was in the **alcohol dehydrogenase (ADH) gene**. This gene encodes for an enzyme that is responsible for converting alcohol to acetaldehyde.

Polymorphisms in this gene can either lead to faster or slower conversion due to increased or decreased enzyme activity respectively. Accordingly, people with different polymorphisms were found to be more likely to experience dizziness, dysphoria or euphoria when drinking alcohol and these side effects of alcohol consumption are known to be important in determining drinking habits. Therefore, polymorphisms in this gene were found to be associated with [alcoholism](#).

Dopamine receptors play an important role in the reward system. Polymorphisms in the dopamine D2 receptor gene (DRD2) were also found to be associated with an increased risk of drug addiction and alcoholism.

Genome-wide association studies also allowed us to understand why certain people are at an increased risk of developing **obesity**, [breast cancer](#) or [Crohn's disease](#) when there is no clear hereditary pattern.

These findings of genome-wide associations with different behavioral traits showed with certainty that genetics are important in determining our behavior.

References

McGue M. [The End of Behavioral Genetics?](#) Xin Li Xue Bao. 2008. 1;40(10):1073-1087. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20072716>

Legal Note: Unless otherwise stated, all rights reserved by Lecturio GmbH. For further legal regulations see our [legal information page](#).